

# Application Protocol - hiPS



## ScreenFect®A-plus Transfection Reagent

### Package Contents

Cat. No.	ScreenFect®A-plus	Dilution Buffer
S-6001-2	0.2 ml	10 ml
S-6001	1.0 ml	50 ml
S-6001-3	5 x 1.0 ml	5 x 50 ml

### Storage Conditions

Store ScreenFect® Reagents at 4 °C. Do not freeze. For optimal long term activity, do not allow ScreenFect® Reagents to warm to room temperature each time it is used.

After several months of storage without using the reagent a slight precipitation might occur. If vortexed thoroughly, this has no influence on the performance of ScreenFect® Reagents.


### General Considerations

For optimal results, amounts of ScreenFect®A-plus and nucleic acid (NA) need to be optimized for each cell type and each NA used. An optimization protocol is provided in our product manual which can be downloaded from our homepage.

We strongly recommend the One-Step transfection method for all of our products. For transfection of adherent cells, remove the used medium and mix fresh medium with the transfection complexes. Then add the mix to the cells.

ScreenFect®A-plus is suitable for pDNA and RNA transfection (view our manual for optimized protocols). For best results in mRNA and siRNA delivery, test our specialized reagents.

## ScreenFect® Protocol: One-Step Transfection



## ScreenFect® Products

### ScreenFect®A

Multipurpose reagent (most suitable for pDNA transfection, suitable for RNA applications) with very low cytotoxicity.

### ScreenFect®A-plus

Multipurpose reagent with optimized formulation requiring less reagent per transfection.

### ScreenFect®siRNA

Specialized reagent for best performance in siRNA delivery.

### ScreenFect®65 + Booster

Reagent kit for protein production in HEK suspension cells.

For additional information regarding ScreenFect®A-plus and other ScreenFect® Products, visit our homepage ([www.screenfect.com](http://www.screenfect.com)) and view our product pages and instruction manuals.

# Application Protocol - hiPS



Protocol for the transfection of **hiPS** cells with pDNA

Component	Procedure for one well (12-well-plate)	12-well
<b>1 Reagent Dilution</b>	Dilute 2 µl of ScreenFect®A-plus in OptiMEM to a final volume of 80 µl and mix thoroughly.  <i>Important: Vortex the reagent once per day of use. Add ScreenFect®A-plus reagent directly into supplied buffer with rapid pipette mixing or vortexing.</i>	2 µl reagent 80 µl dilution
<b>2 pDNA Dilution</b>	Dilute a total of 4 µg pDNA in OptiMEM to a final volume of 80 µl.  <i>Tip: Include a positive control for quick and easy detection of transfection (e.g. using GFP plasmid and fluorescence microscopy).</i>	4 µg 80 µl dilution
<b>3 Complex formation</b>	Combine the diluted ScreenFect®A-plus and DNA and mix immediately using 10 rapid pipette strokes. Leave for 20 min at room temperature for complex formation.  <i>Important: Do not vortex!</i>	160 µl complexes
<b>4 Cell preparation &amp; transfection</b>	Add 740 µl freshly detached and resuspended cells to the complexes and mix with pipette.  <i>Tip: The time-saving reverse cell transfection method may not be suited for all cell types. To transfect adherent cells, first remove and discard medium from cells, then add 740 µl fresh culture medium to transfection complexes, mix with pipette and immediately apply to cells.</i>	Add 740 µl cell suspension
<b>5 Cell plating</b>	Transfer the cells and complexes to one well of a 12-well plate.	Transfer cells with complexes to plate

**Note:** This protocol is a guideline. Values are suitable for easy for transfect cell lines. This protocol does not replace optimization experiments. View our manual for instructions. Serum does not affect the performance of ScreenFect®A-plus but we recommend avoiding antibiotics. Cells must be mycoplasma free, in exponential growth phase and have even plating density across the entire surface area.